

1. feladat

$$a) \iint_T \frac{1}{\sqrt{x^2+y^2+1}} dT \quad T = \{(x,y) \mid x^2+y^2 \leq 1, x,y \geq 0\}$$

Hagyományosán: $x \in [0;1] \quad y \in [0; \sqrt{1-x^2}]$

Polarkoordináta: $x = r \cos \varphi \quad y = r \sin \varphi$

$$\underline{\underline{J}} = \begin{bmatrix} \cos \varphi & -r \sin \varphi \\ \sin \varphi & r \cos \varphi \end{bmatrix} \Rightarrow \det \underline{\underline{J}} = r$$

$$r \in [0;1] \quad \varphi \in [0; \pi/2]$$

$$\int_0^1 \int_0^{\pi/2} \frac{1}{\sqrt{r^2+1}} r d\varphi dr = \frac{\pi}{4} \int_0^1 \underbrace{2r}_{g'} \cdot \underbrace{(r^2+1)^{-1/2}}_{f \circ g} dr =$$

$$= \frac{\pi}{4} \cdot 2 (r^2+1)^{1/2} \Big|_0^1 = \frac{\pi}{2} (\sqrt{2}-1)$$

$$b) f(x,y) = x^2+y^2 \quad T = \{(x,y) \mid (x-3)^2 + (y-2)^2 \leq 1\}$$

$$\begin{cases} x = 3 + r \cos \varphi \\ y = 2 + r \sin \varphi \end{cases} \quad \det \underline{\underline{J}} = r \quad r \in [0;1] \quad \varphi \in [0; 2\pi]$$

$$f(r,\varphi) = (3+r \cos \varphi)^2 + (2+r \sin \varphi)^2 = 13 + 6r \cos \varphi + 4r \sin \varphi + r^2$$

$$\int_0^1 \int_0^{2\pi} 13r + 6r \cos \varphi + 4r \sin \varphi + r^3 d\varphi dr = \int_0^1 26\pi r + 2\pi r^3 dr =$$

$$= 13\pi + \frac{\pi}{2} = \frac{27\pi}{2}$$

$$c) f(x,y) = |2xy| \quad T = \left\{ \frac{x^2}{9} + \frac{y^2}{4} \leq 1 \wedge x, y \geq 0 \right\}$$

$$\begin{cases} x = 3r \cos \varphi & r \in [0, 1] \\ y = 2r \sin \varphi & \varphi \in [0, \pi/2] \end{cases}$$

$$\underline{J} = \begin{bmatrix} 3 \cos \varphi & -3r \sin \varphi \\ 2 \sin \varphi & 2r \cos \varphi \end{bmatrix} \Rightarrow \det \underline{J} = 6r$$

$$\int_0^{\pi/2} \int_0^1 2 \cdot 3r \cos \varphi \cdot 2r \sin \varphi \cdot 6r \, dr \, d\varphi =$$

$$\int_0^{\pi/2} \int_0^1 72r^3 \cos \varphi \sin \varphi \, dr \, d\varphi = \int_0^{\pi/2} \underbrace{18}_{f'} \cos \varphi \underbrace{\sin \varphi}_{f} \, d\varphi =$$

$$= 9 \sin^2 \varphi \Big|_0^{\pi/2} = 9$$

$$d) f(x,y) = 4xy^3 \quad T = \left\{ 1 \leq x^2 + y^2 \leq 4; x \geq 0; y \geq \frac{x}{\sqrt{3}} \right\}$$

$$\begin{cases} x = r \cos \varphi & r \in [1, 2] \\ y = r \sin \varphi & \varphi \in [\pi/6, \pi/2] \end{cases} \quad \det \underline{J} = r$$

$$\int_{\pi/6}^{\pi/2} \int_1^2 4r \cos \varphi \cdot r^3 \sin^3 \varphi \, r \, dr \, d\varphi = \int_{\pi/6}^{\pi/2} \int_1^2 4r^5 \cos \varphi \sin^3 \varphi \, dr \, d\varphi$$

$$= \int_{\pi/6}^{\pi/2} \frac{4}{6} (2^6 - 1) \cos \varphi \sin^3 \varphi \, d\varphi = \int_{\pi/6}^{\pi/2} \underbrace{42}_{f'} \cos \varphi \underbrace{\sin^3 \varphi}_{f^3} \, d\varphi$$

$$= 42 \frac{\sin^4 \varphi}{4} \Big|_{\pi/6}^{\pi/2} = \frac{42}{4} \left(1 - \frac{1}{24} \right) = \frac{21}{2} \cdot \frac{15}{16} = \frac{315}{32}$$

$$e_j) f(x,y) = \sin(x^2+y^2) \quad T = \{x^2+y^2 \leq 2^2; x,y > 0\}$$

$$x = r \cos \varphi \quad r \in [0; 2] \quad \det \underline{\underline{J}} = r$$

$$y = r \sin \varphi \quad \varphi \in [0; \pi/2]$$

$$\int_0^2 \int_0^{\pi/2} \sin(r^2) r d\varphi dr = \int_0^2 \frac{\pi}{2} \sin(r^2) r dr = \frac{\pi}{4} \int_0^2 2r \sin r^2 dr$$

$$= \frac{\pi}{4} (-\cos r^2) \Big|_0^2 = \frac{\pi}{4} (1 - \cos 4)$$

2. feladat

$$a_j) \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} 2^{-x^2-y^2} dx dy$$

$$\int_0^{\infty} \int_0^{2\pi} 2^{-r^2} r d\varphi dr = \int_0^{\infty} (-\pi) (-2r) 2^{-r^2} dr =$$

$$-\pi \cdot 2^{-r^2} \ln 2 \Big|_0^{\infty} = \pi \cdot \ln 2$$

$$b_j) \int_0^{\infty} e^{-x^2} dx \Rightarrow \text{Gauss-görbe}$$

elemi úton nem integrálható!

$$J^2 = \int_0^{\infty} e^{-x^2} dx \cdot \int_0^{\infty} e^{-y^2} dy = \int_0^{\infty} \int_0^{\infty} e^{-x^2-y^2} dx dy =$$

$$\int_0^{\infty} \int_0^{\pi/2} -\frac{1}{2} (2r) e^{-r^2} d\varphi dr = \int_0^{\infty} -\frac{\pi}{4} (2r) e^{-r^2} dr =$$

$$= -\frac{\pi}{4} e^{-r^2} \Big|_0^{\infty} = \frac{\pi}{4} \Rightarrow J = \frac{\sqrt{\pi}}{2}$$

3. feladat

$$V = \int_{x^2+y^2+z^2 \leq R} dV$$

$$\begin{cases} x = r \sin \vartheta \cos \varphi & r \in [0; R] \\ y = r \sin \vartheta \sin \varphi & \vartheta \in [0; \pi] \\ z = r \cos \vartheta & \varphi \in [0; 2\pi] \end{cases}$$

$$dV = r^2 \sin \vartheta \, dr \, d\vartheta \, d\varphi$$

$$\begin{aligned} V &= \int_0^{2\pi} \int_0^{\pi} \int_0^R r^2 \sin \vartheta \, dr \, d\vartheta \, d\varphi = \int_0^{2\pi} \int_0^{\pi} \frac{R^3}{3} \sin \vartheta \, d\vartheta \, d\varphi \\ &= \int_0^{2\pi} \frac{R^3}{3} (-\cos \vartheta) \Big|_0^{\pi} d\varphi = \int_0^{2\pi} \frac{R^3}{3} (1+1) d\varphi = \frac{4R^3}{3} \pi \end{aligned}$$

4. feladat

$$f(x,y,z) = z \sqrt{x^2+y^2} \quad T = \{x^2+y^2 \leq 1; 0 \leq z \leq 2; x,y > 0\}$$

$$\begin{cases} x = r \cos \varphi & r \in [0; 1] \\ y = r \sin \varphi & \varphi \in [0; \pi/2] \\ z = z & z \in [0; 2] \end{cases} \quad \det \underline{J} = r$$

$$\int_0^2 \int_0^1 \int_0^{\pi/2} z r^2 \, d\varphi \, dz \, dr = \frac{\pi}{2} \cdot \frac{1}{3} \cdot \frac{4}{2} = \frac{\pi}{3}$$

5. Seradat

$$f(x,y,z) = \sqrt{x^2+y^2+z^2} \quad T = \{x^2+y^2+z^2 \leq 1\}$$

$$x = r \sin \vartheta \cos \varphi$$

$$y = r \sin \vartheta \sin \varphi$$

$$z = r \cos \vartheta$$

$$r \in [0, 1]$$

$$\vartheta \in [0, \hat{\pi}]$$

$$\varphi \in [0, 2\hat{\pi}]$$

$$\det \underline{J} = r^2 \sin \vartheta$$

$$x^2 + y^2 + z^2 = r^2$$

$$\int_0^{\hat{\pi}} \int_0^{2\hat{\pi}} \int_0^1 r^3 \sin \vartheta \, dr \, d\varphi \, d\vartheta = \int_0^{\hat{\pi}} \frac{1}{4} \cdot 2\hat{\pi} \cdot \sin \vartheta \, d\vartheta =$$

$$= \frac{\hat{\pi}}{2} (-\cos \vartheta) \Big|_0^{\hat{\pi}} = \hat{\pi}$$

6. Seradat

$$\underline{r}(t) = \begin{bmatrix} t^2 + 1 \\ 5t \\ 3t - 2 \end{bmatrix} \quad \underline{v} = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$$

$$\underline{s}(t;s) = \underline{r}(t) + s \underline{v}$$

$$t, s \in \mathbb{R}$$

$$\underline{s}(t;s) = \begin{bmatrix} t^2 + 1 \\ 5t \\ 3t - 2 \end{bmatrix} + s \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$$

7. Seradat

$$\underline{s}(t;s) = \begin{bmatrix} 2 \cos t \\ 2 \sin t \\ s \end{bmatrix} \quad \left. \vphantom{\underline{s}(t;s)} \right\} x^2 + y^2 = 2^2$$

$$t \in [0, 2\hat{\pi}] \\ s \in \mathbb{R}$$

8. Seradat

$$\left. \begin{aligned} x(t) &= 5 + 3 \cos t \\ z(t) &= 3 \sin t \end{aligned} \right\}$$

$$\underline{s}(t;s) = \begin{bmatrix} (5 + 3 \cos t) \cos s \\ (5 + 3 \cos t) \sin s \\ 3 \sin t \end{bmatrix} \quad \left. \vphantom{\underline{s}(t;s)} \right\} \begin{aligned} t &\in [0, 2\hat{\pi}] \\ s &\in [0, 2\hat{\pi}] \end{aligned}$$